


MOBIDERM Technology

MOBIDERM is a medical device comprised of foam cubes encased between two non-woven bandages.

MOBIDERM Technology can be used under a reducing bandage (in its bandage or pad forms), or incorporated into mobilizing garments.

The pressure difference created between the support area of the blocks and the surrounding area stimulates the flow of lymphatic fluid, optimizing drainage efficacy.⁽³⁾



Two sizes of foam blocks adapt to the various needs of the affected area



SMALL BLOCKS

- More effective in highly induced areas
- Better for small morphologies or body areas: pediatric patients, hands/foot, breast, and more.



BIG BLOCKS

- Better adapted along the limb

MOBIDERM autofit garments are recommended for patients with all stages of lymphedema, dysmorphic or regular morphology, progressive edema.

MOBIDERM Standard




Mitten



Glove

MOBIDERM garments are also available in non adjustable version.

ORDERING INFORMATION



*Note: We recommend going up one size to accommodate for upper arm coverage



Size*	Hand Circumference (cA)	Wrist Circumference (cC)	Forearm Circumference (at the widest point) (cD)	Axillary Arm Circumference (cG)	Regular $\ell = 40-45$ cm	Long $\ell = 45-50$ cm
Right	1	17-21 cm	15-19 cm	22-28 cm	25-32 cm	373304 20100262 373304 20100272
	2	17-21 cm	16-20 cm	24-30 cm	29-36 cm	373304 20200262 373304 20200272
	3	18-22 cm	17-21 cm	26-32 cm	33-40 cm	373304 20300262 373304 20300272
	4	18-22 cm	17-21 cm	28-34 cm	37-44 cm	373304 20400262 373304 20400272
	5	19-23 cm	18-22 cm	30-36 cm	41-48 cm	373304 20500262 373304 20500272
	6	20-24 cm	19-23 cm	31-37 cm	45-52 cm	373304 20600262 373304 20600272
Left	1	17-21 cm	15-19 cm	22-28 cm	25-32 cm	373304 20100261 373304 20100271
	2	17-21 cm	16-20 cm	24-30 cm	29-36 cm	373304 20200261 373304 20200271
	3	18-22 cm	17-21 cm	26-32 cm	33-40 cm	373304 20300261 373304 20300271
	4	18-22 cm	17-21 cm	28-34 cm	37-44 cm	373304 20400261 373304 20400271
	5	19-23 cm	18-22 cm	30-36 cm	41-48 cm	373304 20500261 373304 20500271
	6	20-24 cm	19-23 cm	31-37 cm	45-52 cm	373304 20600261 373304 20600271

Size	Wrist Circumference (cC)	Hand Circumference (cA)	Glove		Mitten
			Right	Left	Right/Left
1	14-16 cm	15-17 cm	373201 20100201	373201 20100202	373101 20100299
2	16-18 cm	17-19 cm	373201 20200201	373201 20200202	373101 20200299
3	17-19 cm	19-21 cm	373201 20300201	373201 20300202	373101 20300299
4	19-21 cm	21-23 cm	373201 20400201	373201 20400202	373101 20400299
5	20-22 cm	23-25 cm	373201 20500201	373201 20500202	373101 20500299
6	21-23 cm	25-27 cm	373201 20600201	373201 20600202	373101 20600299

(1) Mestre - Interest of an auto-adjustable night-time compression sleeve (MOBIDERM Autofit) in maintenance phase of upper limb lymphoedema: the MARILYN pilot RCT - Support Care Cancer (25:2455-2462) - 2017 - Page 7

(2) Quéré - Prospective multicenter observational study of lymphoedema therapy: POLIT study - Vascular Disease Journal - 2014 - Pages 4 and 5

(3) Todd - Mobiderm Autofit: an adjustable sleeve that enables patients to self-manage lymphoedema - Chronic oedema - April 2018 - Page 6

(4) Only in following countries: Algeria, Austria, Belgium, Canada, China, Czech Republic, Denmark, Finland, France, Germany, Hungary, India, Morocco, Italy, Netherlands, Poland, Portugal, Romania, Russian Federation, Slovakia, Slovenia, Spain, Sweden, Switzerland, Tunisia, United-Kingdom

Biflex - Indications: Venous insufficiency, simple and complicated varicose veins, reduction of venous and posttraumatic edema, dermatitis / lipodermatosclerosis, varicose ulcers, deep vein thrombosis (prevention and confirmed accident), post-thrombotic syndrome, following sclerotherapy, treatment of lymphedema. Contraindications: PAD (peripheral arterial disease), skin wounds, abscess, infected skin conditions, extra-anatomic bypass, uncontrolled heart failure.

Biflexideal® - Indications: Treatment of lymphedema, simple and complicated varicose veins, following sclerotherapy, following surgery, reduction of post-traumatic edema, varicose ulcers. Contraindications: Do not apply directly to an open wound.

Mobiderm Autofit and Standard - Indications: Maintenance of volume reduction in lymphedema. Contraindications: Skin infection of the limb or acute inflammation, known allergy to components used, septic thrombosis, severe peripheral neuropathy of the limb. Lower limb - PAD (peripheral arterial disease) of lower limbs with ABPI < 0.6, congestive heart failure, advanced diabetic microangiopathy, phlegmasia cerulea dolens, extra-anatomic bypass. Upper limb - pathology of brachial plexus, vasculitis of the extremities.

Availability of these products might vary from a given country or region to another, as a result of specific local regulatory approval or clearance requirements for sale in such country or region. All the medical devices mentioned on this document are CE marked according to the European council directive 93/42/EEC and its relatives or the Regulation 2017/745 on medical devices, unless specifically identified as "NOT CE MARKED". Products mentioned in this document are CE class I devices. Please contact Thuasne should you need any additional information on devices classification. Please read carefully the instructions for use, indications and contraindications of the product. Last revision date: 2020/07. Ref: 002

THERAFIRM®
THUASNE

800-821-3094 | cs@knightrite.com
wholesale.therafirm.com



THUASNE

- Order
- Quotation
- Renewal

MOBIDERM

Patient's First Name: _____

Patient's Last Name: _____

Gender: M F Pediatric

Patient Height: _____

Case No. for renewal: _____

1st Treatment

Date: _____ Quantity: _____

Left Arm Right Arm

Fill out one for each side

MOBIDERM autofit

Comments: _____

**Note: We recommend going up one size to accommodate for upper arm coverage*

G


D

ℓ

C


A

ℓ



Mitten

A



C

Left Hand Right Hand


Fill out one for each side

MOBIDERM

Mitten Glove

Glove

A



C

Comments: _____
